



Sept. 27th, 2006
AIRS Science Team Meeting

Status of the V5 Cloud Cleared and AIRS Only Regressions

Lihang Zhou, Mitch Goldberg, Chris Barnet, Walter Wolf, Xingpin Liu
Fengying Sun, Zhaohui Cheng, Jennifer Wei, Eric Maddy, Murty Divakarla,



Outline

- What's new since V4
- Cloud Cleared Regression Status
- Cloudy Regression (AIRS-Only) Status
- Future Work



What's New Since V4

- Updated Surface Emissivity Coefficient
- Cloudy Regression
 - Generate the coefficient (Feb. 2004)
 - Implemented and tested in NOAA
 - Installed into JPL (July 2005)
 - Generated coef. trained using George's PLR test and delivered to JPL.



What's New Since V4 (2)

➤ Bug Fix:

- Changed the T(p) routine to splice the UARS climatology above 1.5 mb.
- Fixed the bottom temperature and moisture for topography.
- Turned off the AMSU adjustment for total precip. water.



NOAA Surface Regression

- A synthetic regression is used to derive the surface emissivity coefficients
 - Infrared radiance computed from the ECMWF forecast
 - Surface emissivity model
 - Different surface conditions




NOAA Surface Regression V4

- Based on AIRS Level 2 Simulation System (AL2SS) surface model
 - 2 soil types
 - 3 vegetation types
 - Two types of ice: snow and ice
 - Each footprint is a mixture of components based on NDVI and IGBP surface types



Spectra from AL2SS:





Issues with V4:

- Unrealistic snow/ice spectra used for training.
- Lack of short-wave variation in training emissivity.




NOAA Surface Regression V5

- Use the digitized observations of surface emissivity spectra, provided by NESDIS Joint Center for Satellite Data Assimilation (JCSDA).
 - 12 ice/snow spectra
 - 14 vegetation/soil spectra
 - 3 sets of coefficients generated for:
 - Unfrozen land
 - Ocean
 - Snow/Ice




Ice/Snow EMIS Spectra






Land EMIS Spectra






Results: Monthly Averaged Surface Emissivity






Results: Seasonal Characteristics 2005, 980 cm^{-1}





Results: Seasonal Characteristics 2005, 1120 cm^{-1}





Recent On-going NOAA Regression Emissivity Activities:

- Experiment with adding more surface types, such as desert.
- Including more channels (281 channel set).
- Joined CIMSS for inter-comparison of Surface emissivity products.
- Generating monthly emissivity maps which can be used to build the emissivity climatology after careful analysis.




Recent On-going NOAA Regression Emissivity Activities:

- Monthly emissivity maps:
 - 3x3 global grid
 - From Aug 2003 to current
 - Reprocessing capability
 - Maps From Regression:
 - take all FOVs that ok by all steps
 - (ispare(1)=0 and ispare(2)=0)
 - Maps From Physical Retrieval:
 - amplification factor < 1
 - Liquid Water value (liqwat, gm/cm²) < .001
 - NOAA PC Score < 1.



AIRS Derived Monthly Emissivity Climatology (Regression)


(By Fengying Sun) from $3\times 3^\circ$ grids: $11.8\text{ }\mu\text{m}$ (850 cm^{-1})





AIRS Derived Monthly Emissivity Climatology (Retrieval)

(By Fengying Sun) from $3\times 3^\circ$ grids: $11.8\text{ }\mu\text{m}$ (850 cm^{-1})





Cloudy Regression (1)

- Mitch came up the idea to generate regression coefficient from cloud contaminant radiances in early 2004.
- We implemented and tested the cloudy regression in NOAA offline regression system.
- Installed the cloudy regression into the JPL system in July of 2005.
- Used the same thresholds and training days for the training, as we did for the cloud cleared radiance.



Cloudy Regression Coef.(2)

➤ data sets :

- Sept 6 2002, Jan 25 2003, and June 8 2003
- cloudy radiance

➤ filters to the data :

- $\text{diff} \leq 5$. (predict amsu from ahrs - amsu)
 $\text{score} \leq 1.2$
- $\text{delta} < 15$ (for 12 selected channels, abs of difference between sim and obs less than 15).



Cloudy Regression (3)

- Took recommendation from JPL to use the pseudo lapse rate test to train the regression.
- Delivered the re-trained regression coefficient to JPL in this March.




Pseudo Lapse Rate Test:

- Pseudo Lapse Rate Test (#2):
 $PLR = Tb2388 - Tb2387$
 $PLR < \min(5.0, 6 * \cos(lat))$ if $abs(lat) < 60$ and if $topog < 2000m$
- This test will be applied after all retrievals are finished
- If failed, all quality indicators, including Qual_Cloud_OLR, will be set to 2(bad, do not use for data analysis)
- But retrieved values will be kept for later debugging purpose



Comparison of Thresholds




- Test 1(Black)
 - Black curve could be higher if the granule median is higher
- Test 2(Red)

Sung-Yung Lee




PLR Test






AMSU Diff Test





Global non-frozen Ocean, Day 'Clear', AMSU test vs PLR test:





Future Work: Plans for V6

- Continue our experiments on surface emissivity regression:
 - Adding surface type information.
 - Eigenvector regression (Bill Smith).
 - Channel sets.
 - Build a global emissivity climatology.
 - Use of MODIS as first guess.



Future Work: Plans for V6 (2)

- Improve the atmospheric regression for both cloud cleared and cloudy regression:
 - Better training:
 - Generate coef. Based on sondes (T,Q,O3).
 - Remove channels that are bad in certain epochs :
 - Epoch 1 (Before Nov 2003) and epoch 2 coef. Work well.
 - May introduce artifacts to climate studies.
 - In v6 the goal is to establish a common set for all epochs
 - Better QA:
 - identify the problematic area and develop QA schemes accordingly.
 - Extend our AIRS regression techniques to IASI and vise versa.